

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE CTD Observations of the Arctic Ocean Halocline				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Washington, Applied Physics Laboratory, 1013 N.E. 40th Street, Seattle, WA, 98105				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 2	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

CTD OBSERVATIONS OF THE ARCTIC OCEAN HALOCLINE

Dr. Michael Steele
Applied Physics Laboratory, University of Washington
1013 NE 40th St.
Seattle, WA 98105
mas@apl.washington.edu
Voice: 206-543-6586 FAX: 206-543-3521
Award No.: N00014-95-1-0437

LONG TERM GOALS

We seek to identify the extent, morphology, and genesis of the Arctic Ocean halocline as well as surface and Atlantic Water layers.

OBJECTIVES

The objective is to map the extent of various halocline types and compare their properties with older data and with climatologies.

APPROACH

We have used CTD data taken by the US nuclear submarine Cavalla during the spring of 1995. Three types of CTD observations were used: (i) high-frequency underway CTD measurements recorded by an instrument fixed in the ship's sail, (ii) expendable CTD measurements recorded by an experimental XCTD developed by Sippican and launched through the torpedo tubes, and (iii) traditional through-ice CTD profiles obtained after the submarine surfaced through the ice.

ACCOMPLISHMENTS

We have completed the intercalibration of CTD observations taken by the three methods described in "Approach."

SCIENTIFIC RESULTS

A comparison of our observations with data taken in 1993 and 1991 reveals significant trends and frontal shifts through the decade of the 1990's. A comparison with climatology reveals that these changes had just started in 1991, and supports the idea proposed by others of an abrupt transition in the late 1980's. In particular, we have examined changes in the so-called "Cold Halocline Layer" (CHL), a layer of cold, salty water that lies under the surface layer and above the Atlantic Layer in the Arctic Ocean. The climatological significance of the CHL is that it insulates the surface waters (and thus the sea ice) from the relatively warm Atlantic Layer that lies at several hundred meters depth. Without a CHL, heat is available for mixing into the surface layer, and ice melting is the result. An example is the Antarctic sea ice pack, where a CHL is absent and heat from the Atlantic Layer rises up to the surface freely. The result is much thinner ice (less than a meter thick) than in the Arctic Ocean.

Our results show that the area of the deep Arctic Ocean covered by a true CHL in 1995 was actually quite small: essentially, only the Makarov Basin. Comparison with previous data indicate that the Amundsen Basin used to have a CHL at least through 1991; by 1995 it had disappeared from this basin. The result is an unprecedented salinification of surface waters in the Eurasian Basin. We say “unprecedented” because our comparison with a new Russian-American climatology reveals that in 1995 these waters were saltier than the maximum salinities ever recorded in the 40 year history of the climatology.

A paper has been submitted to the Journal of Geophysical Research on this subject (Steele and Boyd, 1997).

IMPACT FOR SCIENCE

Changes in the water mass structure of the upper Arctic Ocean have recently been documented by others. Our study emphasizes that these changes may influence the overlying sea ice cover by removing the insulating CHL and thereby enhancing sea ice melting. This may provide motivation for an initiative to monitor these large-scale changes in the Arctic Ocean.

TRANSITIONS

None.

RELATED PROJECTS

None.

REFERENCES

- Steele, M. and T. Boyd, Retreat of the Cold Halocline Layer in the Arctic Ocean, submitted to *J. Geophys. Res.*, 1997.
- Steele, M., J. Zhang, D. Rothrock, and H. Stern, The force balance of sea ice in a numerical model of the Arctic Ocean, *J. Geophys. Res.*, in press, 1997.